

GROUP	14	HERBICIDE
-------	----	-----------

HEAT™ WG

Water soluble granular herbicide for pre-seed, pre-emergent and chemfallow application for control of broadleaf weeds

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

GUARANTEE: Saflufenacil..... 70%

Warning: Contains the allergen sulfites

REGISTRATION NO. 29368

PEST CONTROL PRODUCTS ACT

CAUTION – SKIN IRRITANT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

NET CONTENTS: 0.25 kg – 100 kg, Bulk

BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, ON L5R 4H1
1-877-371-2273

HEAT™ is a trademark of BASF.

Table of Contents

1.0	GENERAL INFORMATION.....	2
2.0	PRECAUTIONS	2
3.0	FIRST AID.....	3
4.0	ENVIRONMENTAL HAZARDS.....	3
5.0	STORAGE.....	3
6.0	DISPOSAL	4
7.0	NOTICE TO USER	4
8.0	REGISTERED CROPS.....	4
9.0	DIRECTIONS FOR USE.....	5
9.1	CROP USE RATES – HEAT WG	5
9.2	WEEDS CONTROLLED	6
9.3	TANK MIXES	7
9.3.1	HEAT WG + GLYPHOSATE.....	7
9.4	CROP SPECIFIC RECOMMENDATIONS	8
9.4.1	CHICKPEAS AND PEAS (DRIED FIELD).....	8
9.4.2	CEREALS – BARLEY, CANARY SEED, OATS AND WHEAT (spring, durum, winter)	9
9.4.3	SOYBEANS	9
9.4.4	LENTILS (including imazethapyr and imazamox tolerant lentils e.g., Lentil varieties with the CLEARFIELD® TRAIT)	9
9.4.5	CORN (FIELD, SWEET).....	10
9.5	CHEMFALLOW.....	10
10.0	MIXING INSTRUCTIONS	10
11.0	PRE-HARVEST INTERVAL (PHI)	11
12.0	FOLLOW CROPPING.....	12
13.0	SPRAYING INSTRUCTIONS	12
14.0	RESTRICTIONS AND LIMITATIONS.....	12
15.0	RESISTANCE-MANAGEMENT RECOMMENDATIONS	14

1.0 GENERAL INFORMATION

HEAT WG is a water soluble granular herbicide for broadleaf weed control.

HEAT WG is rapidly absorbed by root and foliar uptake; once absorbed it exhibits mobility in plants. **HEAT WG** is a potent inhibitor of protoporphyrinogen oxidase (PPO). Cell membrane damage induced by inhibition of PPO leads to plant death. Susceptible weeds develop injury symptoms within hours of application under active growing conditions; plant death occurs within 3 to 5 days depending upon growing conditions.

HEAT WG is recommended for pre-seed and pre-emergent applications. **HEAT WG** can also be applied in fallow crop lands and post harvest.

HEAT WG does not control grass weeds. **HEAT WG should always be tank mixed with glyphosate** for broad spectrum weed control.

HEAT WG is a broad spectrum weed resistance management tool for activity on a range of broadleaf weeds.

2.0 PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally. Harmful if swallowed.
3. **DO NOT** get in eyes or on skin. Causes eye and skin irritation. Avoid inhalation of vapor or spray.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.
5. Wear a long-sleeved shirt, long pants, coveralls, chemical resistant gloves and shoes plus socks during mixing, loading, clean-up and repair. In addition, wear goggles or face shield during mixing/loading. Applicators must wear long-sleeved shirt, long pants, coveralls and shoes plus socks.

Applicators treating corn fields must wear chemical resistant coveralls over long-sleeved shirt, long pants and chemical resistant gloves during mixing, loading, application, clean-up and repair, and additionally, goggles or a face shield during mixing and loading. Applicators performing mixing, loading, and application must use a closed mixing/loading system.

6. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
7. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

8. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

3.0 FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

4.0 ENVIRONMENTAL HAZARDS

TOXIC to non-target terrestrial plants. Observe buffer zones specified under Section 14 (Restrictions and Limitations).

5.0 STORAGE

1. Store the product in original, tightly-closed container and do not allow water to be introduced into this container.
2. **DO NOT** ship or store the product near food, feed, seed or fertilizers.
3. Store the product in a cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

6.0 DISPOSAL

For Recyclable containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

7.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

8.0 REGISTERED CROPS

HEAT WG is registered for use prior to the following crops as a pre-seed, pre-emergent, pre-plant or pre-plant incorporated application.

Pre-seed or Pre-emergent
• Barley
• Canary seed
• Chickpeas
• Lentils*
• Oats
• Peas (dried field)
• Wheat (spring, winter and durum)
• Corn
• Soybeans*

* Rate restrictions apply. Refer to crop specific section for details.

Chemfallow

9.0 DIRECTIONS FOR USE

9.1 CROP USE RATES – HEAT WG

For all **HEAT WG** solo applications, use MERGE® Adjuvant at 0.5 L/ha.

Pre-seed or Pre-emergent

Crop	Use Rate (g/ha)
Lentils*	26
Soybean**	26
Barley	26 – 71
Canary seed	26 – 71
Chickpea Kabuli	26 – 71
Corn (field, sweet***)	26 – 71
Oats	26 – 71
Peas (dried field)	26 – 71
Wheat (spring, durum, winter)	26 – 71
Chemfallow	26 – 71

* Rate restrictions apply. Do not use rates higher than 26 g/ha or injury could result. See crop specific section for additional details.

** Rate restrictions apply. Do not use rates higher than 26 g/ha or injury could result. See crop specific section for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.

*** Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

Chemfallow

Crop	Use Rate (g/ha)	Application Timing
Chemfallow	26 – 71	Post-emergence in fallow croplands and post harvest.

9.2 WEEDS CONTROLLED

HEAT WG can be applied pre-seed or pre-emergence to the crop to control weeds listed below.

HEAT WG should always be applied in combination with glyphosate for broad spectrum weed control including grassy weeds.

Weeds Controlled For Rapid Burn down (Pre-seed or Pre-emergent)

Use Rate: 26 g/ha

Weed	Maximum Weed Stage
Kochia	15 cm
Canada fleabane	8 leaf
Cleavers	4-whorl stage
Lamb's-quarters	8 leaf
Narrow-leaved hawk's beard	8 cm
Redroot pigweed	8 leaf
Round-leaved mallow	8 leaf
Stinkweed	8 leaf
Volunteer canola (all types including Roundup Ready®)	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

For Rapid Burndown and Suppression of Secondary Weed Flushes (Pre-seed or Pre-emergent)

Use Rate: 71 g/ha

Weed	Maximum Weed Stage
Redroot pigweed	8 leaf
Stinkweed	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

9.3 TANK MIXES

9.3.1 HEAT WG + GLYPHOSATE

For broad spectrum weed control, **HEAT WG** should always be tank mixed with glyphosate present as isopropylamine salt, di-ammonium salt or potassium salt. **HEAT WG** is compatible with all liquid glyphosate formulations, in which glyphosate is present as isopropylamine salt, di-ammonium salt or potassium salt, as a tank-mix partner.

Tank mixing **HEAT WG** with glyphosate will provide control of all weeds controlled by glyphosate in addition to the broadleaf weeds listed on **HEAT WG** label. Consult the glyphosate label for a complete list of weeds controlled by glyphosate.

Tank mixing is a recognized strategy to delay herbicide resistance as well as improve the weed spectrum controlled. See Section 15.

TANK MIX CROP USE RATES

HEAT WG + Glyphosate Use Rate (360 g/L equivalent) 1.25 – 2.5 L/ha

Crop	HEAT WG Use Rate (g/ha)
Lentils*	26
Soybean**	26
Barley	26 - 71
Canary seed	26 - 71
Chickpea Kabuli	26 - 71
Corn (field, sweet***)	26 - 71
Oats	26 - 71
Peas (dried field)	26 - 71
Wheat (spring, durum, winter)	26 - 71
Chemfallow	26 - 71
* Rate restrictions apply. Do not use rates higher than 26 g/ha or crop injury may result. See crop specific section for additional details.	
** Rate restrictions apply. Do not use rates higher than 26 g/ha or crop injury may result. See crop specific section for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.	
*** Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.	

For tank mix applications of **HEAT WG** with glyphosate, use Merge® Adjuvant or Amigo® at 0.5 L/ha.

Always refer to the tank mix partner herbicide label for precautions, use instructions and crop rotation restrictions. Do not apply tank mix combinations by air.

When a tank mix is used, consult the labels of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarse spray (ASAE) category indicated on the labels for those tank mix partners.

TANK MIX WEEDS CONTROLLED

Weeds Controlled for Rapid Burn Down (Pre-seed or Pre-emergent)

HEAT WG when tank mixed with glyphosate will provide rapid burn down of the following weeds in addition to weeds listed for treatment of **HEAT WG**.

Use Rate: 26 g/ha

Weed	Maximum Weed Stage
Dandelion (top growth burndown control only)	15 cm
Flixweed	8 leaf

9.4 CROP SPECIFIC RECOMMENDATIONS

9.4.1 CHICKPEAS AND PEAS (DRIED FIELD)

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of HEAT WG + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or Amigo at a rate of 0.5 L/ha.
Weeds Controlled	See Section 9.2 for a complete list of weeds controlled by HEAT WG . Refer to the glyphosate label for weeds controlled in addition to those controlled by HEAT WG .
Remarks	Do not apply any other soil applied herbicide with, before, or after applications of HEAT WG + glyphosate. Sufficient data to support such a use pattern is not available and injury could result.

**9.4.2 CEREALS – BARLEY, CANARY SEED, OATS AND WHEAT
(spring, durum, winter)**

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of HEAT WG + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or Amigo at a rate of 0.5 L/ha
Weeds Controlled	See Section 9.2 for a complete list of weeds controlled by HEAT WG . Refer to the glyphosate label for weeds controlled in addition to those controlled by HEAT WG .

9.4.3 SOYBEANS

Timing	Pre-seed or Pre-emergent
Rate	26 g/ha of HEAT WG + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or Amigo at a rate of 0.5 L/ha.
Weeds Controlled	See Section 9.2 for a complete list of weeds controlled by HEAT WG . Refer to the glyphosate label for weeds controlled in addition to those controlled by HEAT WG .
Remarks	Some soybean cultivars may be more sensitive to saflufenacil and injury might occur.

9.4.4 LENTILS (including imazethapyr and imazamox tolerant lentils e.g., Lentil varieties with the CLEARFIELD® trait)

Timing	Pre-seed or Pre-emergent
Rate	26 g/ha of HEAT WG + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 - 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or Amigo at a rate of 0.5 L/ha.
Weeds Controlled	See Section 9.2 for a complete list of weeds controlled by HEAT WG . Refer to the glyphosate label for weeds controlled in addition to those controlled by HEAT WG .

Remarks	<p>Do not apply any other soil applied herbicide with, before, or after applications of HEAT WG + glyphosate. The addition of other soil applied herbicides can increase the sensitivity of lentils to HEAT WG and injury could result.</p> <p>Rainfall shortly after product application can result in slight injury to the crop. Lentils will be more susceptible to injury on coarse texture and low organic matter soils. Injury will usually appear as leaf tissue necrosis on the outer edges of the leaves. Lentils will grow out of injury symptoms, and yield will not be impacted at recommended rates.</p>
---------	--

9.4.5 CORN (field, sweet)

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of HEAT WG + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or Amigo at a rate of 0.5 L/ha.
Weeds Controlled	See Section 9.2 for a complete list of weeds controlled by HEAT WG . Refer to the glyphosate label for weeds controlled in addition to those controlled by HEAT WG .
Remarks	Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

9.5 CHEMFALLOW

Timing	Chemfallow
Rate	26 – 71 g/ha of HEAT WG + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or Amigo at a rate of 0.5 L/ha.
Weeds Controlled	See Section 9.2 for a complete list of weeds controlled by HEAT WG . Refer to the glyphosate label for weeds controlled in addition to those controlled by HEAT WG .
Remarks	Apply to actively growing weeds less than 15 cm in height. Better coverage of the product results in enhanced control of weeds. For application to larger weeds or dense weed infestations, use minimum water volume of 100 L per hectare.

10.0 MIXING INSTRUCTIONS

1. When using **HEAT WG**, always start with a clean sprayer. Thoroughly clean the sprayer by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.

2. Fill clean spray tank half full with clean water. Start agitation system.
3. Using a calibrated measuring device, add the correct amount of **HEAT WG**. Continue to agitate.
4. **For tank mix with glyphosate**, add **HEAT WG** followed by recommended amount of glyphosate.
5. If recommended, add the correct amount of MERGE Adjuvant or Amigo to the tank last.
6. Continue agitation while filling the remainder of the tank with water necessary to fill the spray tank.
7. Continue to agitate or run the by-pass system.
8. After any break in spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit over night.
9. If an oil film starts to build up in the tank, drain it and clean the tank with strong detergent solution.
10. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

Dispose of all rinsings in accordance with provincial regulations.

11.0 PRE-HARVEST INTERVAL (PHI)

The following pre-harvest intervals should be observed for respective crops.

Crop	Pre-Harvest Interval (PHI) in days
Barley	60
Canary seed	60
Chickpea	60
Corn (field, sweet)	60
Lentils	60
Oats	60
Peas (dried field)	60
Soybean	60
Wheat (spring, winter, durum)	60

12.0 FOLLOW CROPPING

These crops can be safely grown safely after an application of **HEAT WG**.

Plant Back Crops In case of crop failure, the following crops can be planted in the same season ¹	Rotational Crops The following crops can be planted anytime in the following season
Barley Canary seed Chickpeas Corn (field and sweet) Lentils* Oats Dry field peas Soybean* Wheat (spring, winter & durum)	Barley Canary seed Canola Chickpeas Corn (field and sweet) Dry beans Flax Lentils Mustard Oats Dry field peas Soybean Wheat (spring, winter & durum)
* Rate restrictions apply. Lentils and soybeans can only be grown as plant back crops provided that a maximum product use rate of 26 g/ha was used in the previous crop.	

¹ A second application of **HEAT WG** cannot be made in the rescue crop.

13.0 SPRAYING INSTRUCTIONS

Water volume and spray pressure

Conventional ground application

Use sprayers equipped with standard flat nozzles. The use of 80°-110° stainless steel flat fan nozzle is recommended for optimum spray coverage with nozzles tilted 45° forward to ensure better coverage.

Thoroughly clean all screens to prevent nozzle clogging. Apply in a water volume of 50-100 L/ha and at a pressure of 240 kPa. For applications to dense weed infestations and thick canopies, use a higher water volume at pressures of 275 kPa.

Better coverage of the product results in enhanced control of weeds.

14.0 RESTRICTIONS AND LIMITATIONS

1. Wash sprayer thoroughly after use to avoid damage to the next crop sprayed.
2. DO NOT mix or apply **HEAT WG** with any other additive, pesticide or with fertilizer except as specifically recommended on this label.
3. DO NOT APPLY BY AIR.
4. DO NOT enter or allow worker entry into treated areas for 12 hours after application.

5. Field corn – Corn forage and silage can be harvested, used as feed or grazed 60 or more days after application of **HEAT WG**.
6. Legume forage (chickpeas, field peas and lentils) may be used as feed or grazed 60 or more days after application of **HEAT WG**.
7. Small grains (wheat, barley and oats) – forage and hay can be used as feed or grazed 30 or more days after application of **HEAT WG**.
8. Soybeans - may be used as feed or grazed 60 or more days after application of **HEAT WG**.
9. DO NOT apply directly to water. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
10. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
11. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
12. DO NOT apply during periods of dead calm. Avoid application when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Buffer zones

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

Method of application	Crop	Buffer zones (metres) required for the protection of terrestrial habitats
Field sprayer*	Lentils, soybean	4
	Barley, canary seed, oats, peas, wheat, corn, chickpea	10

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labeled buffer zone can be reduced by 30%.

15.0 RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **HEAT WG** is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to **HEAT WG** and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **HEAT WG** or other Group 14 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273 or at www.agsolutions.ca.

®All other products listed are registered trademarks of their respective companies.

GROUP	14	HERBICIDE
-------	----	-----------

HEAT™ WG

Water soluble granular herbicide for pre-seed, pre-emergent and chemfallow application for control of broadleaf weeds

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

GUARANTEE: Saflufenacil..... 70%

Warning: Contains the allergen sulfites

REGISTRATION NO. 29368

PEST CONTROL PRODUCTS ACT

CAUTION – SKIN IRRITANT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

NET CONTENTS: 844 g

BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, ON L5R 4H1
1-877-371-2273

HEAT™ is a trademark of BASF.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally. Harmful if swallowed.
3. **DO NOT** get in eyes or on skin. Causes eye and skin irritation. Avoid inhalation of vapor or spray.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.
5. Wear a long-sleeved shirt, long pants, coveralls, chemical resistant gloves and shoes plus socks during mixing, loading, clean-up and repair. In addition, wear goggles or face shield during mixing/loading. Applicators must wear long-sleeved shirt, long pants, coveralls and shoes plus socks.

Applicators treating corn fields must wear chemical resistant coveralls over long-sleeved shirt, long pants and chemical resistant gloves during mixing, loading, application, clean-up and repair, and additionally, goggles or a face shield during mixing and loading. Applicators performing mixing, loading, and application must use a closed mixing/loading system.

6. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
7. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.
8. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL HAZARDS

TOXIC to non-target terrestrial plants.

STORAGE

1. Store the product in original, tightly-closed container and do not allow water to be introduced into this container.
2. **DO NOT** ship or store the product near food, feed, seed or fertilizers.
3. Store the product in a cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.