Approved booklet (Dry Edible Beans [2009-4921]), March 23, 2010

GROUP	2	HERBICIDE

SOLO® WDG HERBICIDE

Water Dispersible Granular Herbicide
in water-soluble bags

AGRICULTURAL

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER AREA OF BRITISH COLUMBIA ONLY

GUARANTEE: Imazamox …….. 70% a.e.

Warning, contains the allergen sulfites

REGISTRATION NO. 25496 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 470 g (4 x 117.5 g water-soluble bags)

WARNING - EYE IRRITANT

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673

READ THE LABEL AND THIS BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

BASF Canada Inc.
100 Milverton Drive
5th Floor
Mississauga, Ontario
L5R 4H1

1-877-371-2273

SOLO® is a registered trademark of BASF.

©BASF Canada Inc., 1998

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY BY AIR.

Skin irritant.

WARNING - EYE IRRITANT

May cause eye damage.

Wear a long-sleeved shirt and long pants during all activities, and wear chemical-resistant gloves and goggles or face shield during mixing/loading, clean-up and repair activities.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.

3. Make the empty container unsuitable for further use.

4. Dispose of the container in accordance with provincial requirements.

5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

ENVIRONMENTAL HAZARDS

This product is highly toxic to non-target plants. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay). Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

FOR FIELD PEAS, IMAZETHAPYR AND IMAZAMOX TOLERANT CANOLA[1] (e.g., Canola varieties with the CLEARFIELD® trait), IMAZETHAPYR AND IMAZAMOX TOLERANT LENTILS[1] (e.g., Lentil varieties with the CLEARFIELD® trait), IMAZAMOX TOLERANT SUNFLOWERS[1] (e.g., Sunflower varieties with the CLEARFIELD® trait) AND IMAZETHAPYR AND IMAZAMOX TOLERANT CANOLA[1] QUALITY *BRASSICA JUNCEA* (e.g. Canola Quality *Brassica juncea* varieties with the CLEARFIELD® trait)

DIRECTIONS FOR USE (Prairie Provinces and Peace River Area of British Columbia only)

SOLO WDG Herbicide is a selective herbicide that can be applied as an early post-emergence treatment in field peas, imazethapyr and imazamox tolerant canola, imazethapyr and imazamox tolerant lentils and imazethapyr and imazamox tolerant Canola Quality *Brassica juncea* (e.g. Canola Quality *Brassica juncea* varieties with the CLEARFIELD trait). When SOLO WDG herbicide is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

Apply using ground equipment only. DO NOT APPLY BY AIR.

DO NOT apply more than once per year.

DO NOT graze the treated field peas, imazethapyr and imazamox tolerant canola, imazethapyr and imazamox tolerant lentils or imazethapyr and imazamox tolerant Canola Quality *Brassica juncea* (e.g. Canola Quality *Brassica juncea*) or cut for hay within 20 days of application.

DO NOT graze treated sunflower plants or cut for straw; sufficient data are not available to support such use.

Pre-harvest Interval: After 60 days, canola, dry edible beans (*Phaseolus vulgaris*), field peas, imazethapyr and imazamox tolerant lentils (e.g., Lentil varieties with the CLEARFIELD trait) and Canola Quality *Brassica juncea* can be harvested. Sunflowers can be harvested after 70 days.

Overspray or drift to sensitive habitats should be avoided. A buffer zone of 11 metres is required between the downwind point of direct application and the closest edge of (1) sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands, and shrublands), (2) sensitive aquatic habitats (such as lakes, rivers, sloughs, ponds, coulees, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands) and (3) estuarine or marine habitats.

DO NOT apply during periods of dead calm or when winds are gusty.

When a tank mixture is used, consult the label of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, coulees, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

BROADLEAF WEED AND GRASS CONTROL

SOLO WDG herbicide at 21 g/ha (15 g a.e./ha) will control only:	
Lamb's-quarters* Stinkweed Wild mustard	Volunteer spring wheat (non-imazamox tolerant wheat)
SOLO WDG herbicide at 29 g/ha (20 g a.e./ha) will control:	
Barnyard grass Cleavers* Cow cockle Green foxtail Green smartweed Japanese Brome Grass* Kochia* Lamb's-quarters Persian darnel Redroot pigweed Shepherd's purse Stinkweed	Volunteer barley Volunteer canary seed Volunteer canola (non-imazethapyr and non-imazamox tolerant canola only) Volunteer durum wheat Volunteer spring wheat (non-imazamox tolerant wheat) Volunteer tame oats Wild buckwheat* Wild mustard Wild oats Yellow foxtail

* = suppression only

Apply in 100 L/ha of water with MERGE® Adjuvant at a rate of 0.5% v/v (e.g. 5 L of MERGE Adjuvant per 1000 L spray solution).

FOR IMAZETHAPYR AND IMAZAMOX TOLERANT CANOLA[1]
(e.g., Canola varieties with the CLEARFIELD trait)

Application should be made from the 2 to 6 true leaf stage of the canola crop and after the weeds have emerged. To control broadleaf weeds, apply SOLO WDG herbicide from the cotyledon to the 4-leaf stage. To control grasses, apply SOLO WDG herbicide from the 1 to 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

FOR IMAZETHAPYR AND IMAZAMOX TOLERANT CANOLA[1] QUALITY *BRASSICA JUNCEA* (e.g. Canola Quality *Brassica juncea* varieties with the CLEARFIELD trait)

Application should be from the 2 to 6 true leaf stage of the crop and after the weeds have emerged. To control broadleaf weeds, apply SOLO WDG herbicide from the cotyledon to the 4-leaf stage. To control grasses, apply SOLO WDG herbicide from the 1 to 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

FOR IMAZAMOX TOLERANT SUNFLOWERS[1]
(e.g., Sunflower varieties with the CLEARFIELD trait):

Application should be made from the 2 to 8 leaf stage of the sunflower crop and after the weeds have emerged. To control broadleaf weeds, apply SOLO WDG herbicide from the cotyledon up to the 4-leaf stage. To control grasses, apply SOLO WDG herbicide from the 1 to 4 true leaf stage up until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

Apply in 100 L/ha of water with MERGE® Adjuvant at a rate of 0.5% v/v (e.g. 5 L of MERGE Adjuvant per 1000 L spray solution).

FOR FIELD PEAS

Application should be made from the 1 to 6 true leaf stage and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 to 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

FOR IMAZETHAPYR AND IMAZAMOX TOLERANT LENTILS[1]
(e.g., Lentil varieties with the CLEARFIELD trait)

Application should be made from the 2 to 6 leaf stage of the CLEARFIELD lentil crop and after the weeds have emerged. To control broadleaf weeds, apply SOLO WDG herbicide from the cotyledon up to the 4-leaf stage. To control grasses, apply SOLO WDG herbicide from the 1 to 4 true leaf stage up until early tillering.

TANK-MIX COMBINATION

SOLO WDG Herbicide + Basagran Forte Liquid Herbicide in Field Peas

To provide broad-spectrum weed control, SOLO WDG herbicide may be tank-mixed with Basagran Forte Liquid Herbicide in field peas.

Follow all recommendations, precautions and restrictions on the tank-mix partner label.

Timing	Early post-emergence
Rate	29 g/ha SOLO WDG herbicide + 894 mL/ha Basagran Forte Liquid Herbicide + 2 L/ha nitrogen source** (UAN 28%)
Water Volume	100 L/ha
Weeds Controlled	This tank-mix combination will provide enhanced weed control vs. SOLO WDG herbicide alone, as well as control of weeds listed on SOLO WDG Herbicide label. Refer to the Broadleaf Weed and Grass Control Section of this label. **Control** / **Suppression** Volunteer CLEARFIELD canola / Wild buckwheat Wild mustard (Group 2 resistant) /
Remark	Application should be made from the 3 - 6 node stage of field peas and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield. For tank mixing, see mixing instructions.

**A reduction in grass control can be observed without the addition of nitrogen source (UAN 28%).

SOLO WDG Herbicide + Basagran Forte Liquid Herbicide in Dry Edible Beans (all types of the species *Phaseolus vulgaris)**

To provide broad-spectrum weed control and to reduce the potential for crop injury SOLO WDG herbicide must be tank-mixed with Basagran Forte Liquid Herbicide in dry edible beans.

Timing	Early post-emergence
Rate	0.029 kg/ha SOLO WDG herbicide + 1.25 L/ha Basagran Forte Liquid Herbicide + 2 L/ha nitrogen source** (UAN 28%)
Water Volume	100 L/ha
Weeds Controlled	This tank-mix combination will provide enhanced weed control vs. SOLO WDG herbicide alone, as well as control of weeds listed on SOLO WDG Herbicide label. Refer to the Broadleaf Weed and Grass Control Section of this label. **Control**: Group 2 resistant wild mustard; Stork's bill; Volunteer canola (including biotypes with the CLEARFIELD [CL], ROUNDUP READY and LIBERTY LINK traits) **Suppression**: Prostrate pigweed; Spiny annual sow thistle; Wild buckwheat **Nitrogen source: A reduction in grass control can be observed without the addition of nitrogen source (UAN 28%).
Remark	Solo WDG Herbicide is only registered for use on dry beans when tank-mixed with Basagran Forte Liquid Herbicide + UAN. Application should be made after 1st trifoliate leaf has fully expanded up to the 2nd trifoliate leaf stage of the dry bean and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield. For tank mixing, see mixing instructions. Follow all recommendations, precautions and restrictions on the tank-mix partner label. Do not harvest the treated crop within 60 days of application.
***Warning**	Only the following dry bean types have been tested and demonstrate acceptable tolerance: Pinto, pink, red mexican, cranberry, black, great northern and navy. Since not all types of dry beans of the species *Phaseolus vulgaris* have been tested, first application should be limited to a small area or consult your seed supplier or BASF Business Representative for more information.

NOTE TO USER

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than BASF Canada and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. BASF Canada itself makes no representation or warranty with respect to performance (efficacy) and crop tolerance (phytotoxicity) claims for this product when used on the crops listed above.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerances arising, and agree to hold BASF Canada harmless from any claims based on efficacy or phytotoxicity in connection with the uses described above.

Dry common bean varieties may vary in their tolerance to herbicides, including to **SOLO WDG Herbicide + Basagran Forte Liquid Herbicide** tank mix. In particular, white (navy) beans are more susceptible to herbicide injury which can result in delayed maturity. Since not all dry common bean varieties have been tested for tolerance to **SOLO WDG Herbicide + Basagran Forte Liquid Herbicide** tank mix, first use of **this tank mix** should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice.

SOLO WDG Herbicide + Lontrel Dry Soluble Granular Herbicide in imazethapyr and imazamox tolerant canola (e.g. Canola varieties with the CLEARFIELD trait)

SOLO WDG herbicide may be tank-mixed with Lontrel Dry soluble granular herbicide in imazethapyr and imazamox tolerant canola (e.g. canola varieties with the CLEARFIELD trait).

Timing	Early post-emergence
Rate	29 g/ha SOLO WDG Herbicide + 100 g/ha Lontrel Dry Granular Herbicide
Water Volume	100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant – 0.5% v/v
Weeds Controlled	This tank-mix combination will provide control of weeds as listed on the SOLO WDG Herbicide. Refer to Broadleaf weed and Grass control Section. Improved weed control with the tank mix vs. SOLO WDG Herbicide alone. **Control** Wild buckwheat
Remark	Application should be made from the 2 - 6 leaf stage of the canola crop and after the weeds have emerged. To control broadleaf weeds, apply SOLO WDG herbicide from the cotyledon to the 4-leaf stage. To control grasses, apply SOLO WDG herbicide from the 1 - 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield. For tank mixing, see mixing instructions. Refer to Tank mix rotational crop restrictions for details.

Tank Mix Rotational Crop Restrictions
SOLO WDG Herbicide + Lontrel Dry Soluble Granular Herbicide in imazethapyr and imazamox tolerant canola (e.g. Canola varieties with the CLEARFIELD trait)

Fields previously treated with tank mix can be seeded to wheat, barley, oats and rye (not under-seeded to forage legumes, clover or alfalfa), canola, corn, field peas, flax, forage grasses or can be summer-fallowed. Seed only those crops listed above in the year following treatment.

Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application, delay seeding field peas an additional 12 months (total 22 months following application).

Contact your local BASF Canada representative or retailer for more information before seeding field peas following drought conditions in the previous year.

Tank Mix Pre-harvest / Grazing Interval
Do not harvest the treated crop within 60 days of application.

MIXING INSTRUCTIONS

1. Use 100 L/ha of water.
2. Use a 50-mesh (or coarser) filter screen.
3. Fill the spray tank with three-quarters of the required amount of clean water, start agitation and continue agitation throughout the entire mixing and spraying procedure.
4. Add the required amount of SOLO WDG herbicide soluble bag(s) directly into the sprayer through the tank opening.
5. For SOLO WDG Herbicide+ Basagran Forte liquid herbicide tank mix, add Basagran Forte Liquid Herbicide followed by recommended amount of nitrogen source (UAN 28% or ammonium sulphate).
6. For SOLO WDG Herbicide + Lontrel Dry soluble granular herbicide tank mix, add recommended Lontrel Dry soluble granular herbicide.
7. Agitate for at least ten minutes to dissolve the herbicide. After the herbicide is dissolved, continue agitation.
8. If required, add the required amount of MERGE Adjuvant. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. HALT®).
9. **DO NOT** use MERGE Adjuvant for SOLO WDG herbicide + Basagran Forte Liquid Herbicide tank mix.
10. Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again.
11. Between loads of SOLO WDG Herbicide, Basagran Forte Liquid Herbicide or Lontrel Dry soluble granular herbicide, check in-line and nozzle screens and rinse clean if necessary.
12. Upon completion of spraying SOLO WDG herbicide or the tank mix, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

FOLLOW CROPPING

Initial crop injury to non-imazethapyr and non-imazamox tolerant canola (non-CLEARFIELD canola) may be observed. Avoid spray overlap as yield reduction may result. The following crops may be grown safely the year following an application:

Canary seed
Field corn
Field peas
Imazethapyr and imazamox tolerant canola[1] (CLEARFIELD Canola)
Non-imazethapyr and non-imazamox tolerant canola (non-CLEARFIELD canola)
Lentils
Spring wheat
Durum wheat
Spring barley
Sunflower
Tame oats
Flax
Chickpeas
Imazamox tolerant sunflowers (CLEARFIELD Sunflowers)

The following crop may be grown safely two years following an application:

Mustard (condiment type only)

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, SOLO WDG herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to SOLO WDG herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of SOLO WDG herbicide or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273 or at www.agsolutions.ca.

CLEARFIELD® is a registered trademark of BASF.
MERGE® is a registered trademark of BASF Canada Inc.
®All other products listed are registered trademarks of their respective companies.

[1]*Methods of growing plants which are resistant to certain acetohydroxyacid synthase (AHAS) herbicides are protected by Canadian Patent No. 1341465. The use of this product in the practice of patented methods could constitute patent infringement. The purchase of this product conveys no license to the purchaser to practice the patented methods. Further information can be obtained by contacting BASF Canada at 1-877-371-2273.*

Approved container (Dry Edible Beans [2009-4921]), March 23, 2010

GROUP	2	HERBICIDE

SOLO® WDG HERBICIDE

Water Dispersible Granular Herbicide
in water-soluble bags

AGRICULTURAL

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER AREA OF BRITISH COLUMBIA ONLY

GUARANTEE: Imazamox …….. 70% a.e.

Warning, contains the allergen sulfites

REGISTRATION NO. 25496 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 470 g (4 x 117.5 g water-soluble bags)

WARNING - EYE IRRITANT

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

BASF Canada Inc.
100 Milverton Drive
5th Floor
Mississauga, Ontario
L5R 4H1

1-877-371-2273

SOLO® is a registered trademark of BASF.

©BASF Canada Inc., 1998

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY BY AIR.

Skin irritant.

WARNING - EYE IRRITANT

May cause eye damage.

Wear a long-sleeved shirt and long pants during all activities, and wear chemical-resistant gloves and goggles or face shield during mixing/loading, clean-up and repair activities.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.

3. Make the empty container unsuitable for further use.

4. Dispose of the container in accordance with provincial requirements.

5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Approved pouch label (Dry Edible Beans [2009-4921]), March 23, 2010

SOLO WDG HERBICIDE

70% Water Dispersible Granular Herbicide

AGRICULTURAL

NET CONTENTS: 117.5 g

GUARANTEE: Imazamox … 70% active (a.e.)

Warning, contains the allergen sulfites

WARNING - EYE IRRITANT

KEEP OUT OF REACH OF CHILDREN